



TITLE:

Three dimensional hypersurface purely elliptic singularities

AUTHOR(S):

WATANABE, KIMIO

CITATION:

WATANABE, KIMIO. Three dimensional hypersurface purely elliptic singularities. 数理解析研究所講究録 1998, 1037: 69-73

ISSUE DATE:

1998-04

URL:

<http://hdl.handle.net/2433/61951>

RIGHT:

Three dimensional hypersurface purely elliptic singularities

KIMIO WATANABE*

Abstract

Three dimensional hypersurface purely elliptic singularities are classified into three classes according to the shape of their Newton boundaries.

A lot of examples of their defining equations are obtained from the defining equations of hypersurface simple $K3$ singularities. There are at least 95 types for the defining equations of hypersurface purely elliptic singularities of the type $(0, 1)$ or $(0, 0)$.

1 Introduction

In the theory of normal two-dimensional singularities, simple elliptic singularities and cusp singularities are regarded as the most reasonable class of singularities after rational double points. They are characterized as two-dimensional purely elliptic singularities of $(0,1)$ -type and of $(0,0)$ -type, respectively. What are natural generalizations in three-dimensional case of simple elliptic singularities. The notion of a simple $K3$ singularity was defined in [4] as a three-dimensional isolated Gorenstein purely elliptic singularity of $(0,2)$ -type. A simple $K3$ singularity is characterized as a normal three-dimensional isolated singularity such that the exceptional set of any \mathbf{Q} -factorial terminal modification is a three-dimensional $K3$ surface (see [4]). Here we are interested in three-dimensional hypersurface purely elliptic singularities of $(0, i)$ -type for $i = 0$ or $i = 1$. Let $f \in \mathbf{C}[z_0, z_1, z_2, z_3]$ be a polynomial which

*Partly supported by the Grant-in-Aid for Co-operative Research as well as Scientific Research, the Ministry of Education, Science and culture, Japan.

1991 *Mathematics subject Classification*. Primary 14M25; Secondary 14J10, 52B20, 33C80.

is non-degenerate with respect to its Newton boundary $\Gamma(f)$ in the sense of [5], and whose zero locus $X = \{f = 0\}$ in \mathbb{C}^4 has an isolated singularity at the origin $0 \in \mathbb{C}^4$. Then the condition for the singularity (X, x) to be a purely elliptic singularity of $(0, 0)$ -type is given by a property of the Newton boundary of $\Gamma(f)$ of f .

In this paper, we give the method to obtain the principal parts of defining equations, which define three-dimensional hypersurface purely elliptic singularities of $(0, i)$ -type for $i = 0$ to $i = 1$.

2 Preliminaries

In this section, we recall some definitions and results from [1], [4] and [6].

First we define the plurigenera $\delta_m, m \in \mathbb{N}$, for normal isolated singularities and define purely elliptic singularities. Let (X, x) be a normal isolated singularity in an n -dimensional analytic space X , and $\pi : (M, E) \rightarrow (X, x)$ a good resolution. In the following, we assume that X is a sufficiently small Stein neighbourhood of x .

DEFINITION. ([6]) Let (X, x) be a normal isolated singularity. For any positive integer m ,

$$\delta_m(X, x) := \dim_{\mathbb{C}} \Gamma(X - \{x\}, \mathcal{O}(mK)) / L^{2/m}(X - \{x\}),$$

where K is the canonical line bundle on $X - \{x\}$.

Then δ_m is finite and does not depend on the choice of a Stein neighborhood.

DEFINITION. ([6]) A singularity (X, x) is said to be purely elliptic if $\delta_m = 1$ for every $m \in \mathbb{N}$.

When X is a two-dimensional analytic space, purely elliptic singularities are quasi-Gorenstein singularities, i.e., there is a nowhere vanishing holomorphic 2-form on $X - \{x\}$ (see [2]). But in higher dimension, purely elliptic singularities are not always quasi-Gorenstein (see [3]).

In the following, we assume that (X, x) is quasi-Gorenstein. Let $E = \bigcup E_i$ be the decomposition of the exceptional set E into its irreducible components, and write

$$K_M = \pi^* K_X + \sum_{i \in I} m_i E_i - \sum_{j \in J} m_j E_j$$

with $m_i \geq 0, m_j \geq 0$. Ishii [1] defined the essential part of the exceptional set E as $E_j = \sum_{j \in J} m_j E_j$, and showed that if (X, x) is purely elliptic, then $m_j = 1$ for all $j \in J$.

DEFINITION. ([1],[6]) A quasi-Gorenstein purely elliptic singularity (X, x) is of $(0, i)$ -type if $H^{n-1}(E_J, \mathcal{O}_E)$ consists of the $(0, i)$ -Hodge component $H^{0, i}(E_J)$, where

$$C \simeq H^{n-1}(E_J, \mathcal{O}_E) = Gr_F^0 H^{n-1}(E_J) = \bigoplus_{i=1}^{n-1} H^{0, i}(E_J)$$

n -dimensional quasi-Gorenstein purely elliptic singularities are classified into $2n$ classes, including the condition that the singularity is Cohen-Macaulay or not.

Next we consider the case where (X, x) is a hypersurface singularity defined by a non-degenerate polynomial $f = \sum a_\nu z^\nu \in \mathbb{C}[z_0, z_1, \dots, z_n]$, and $x = 0 \in \mathbb{C}^{n+1}$. Recall that the Newton boundary $\Gamma(f)$ of f is the union of the compact faces of $\Gamma_+(f)$, where $\Gamma_+(f)$ is the convex hull of $\bigcup_{a_\nu \neq 0} (\nu + \mathbb{R}_0^{n+1})$ in \mathbb{R}^{n+1} . For any face Δ of $\Gamma_+(f)$, set $f_\Delta := \sum_{\nu \in \Delta} a_\nu z^\nu$. We say f to be nondegenerate, if

$$\frac{\partial f_\Delta}{\partial z_0} = \frac{\partial f_\Delta}{\partial z_1} = \dots = \frac{\partial f_\Delta}{\partial z_n} = 0$$

has no solution in $(\mathbb{C}^*)^{n+1}$ for any face Δ . Where f is nondegenerate, the condition for (X, x) to be a purely elliptic singularity of $(0, i)$ -type is given as follows:

THEOREM 2.1 *Let f be a nondegenerate polynomial and suppose $X = \{f = 0\}$ has an isolated singularity at $x = 0 \in \mathbb{C}^{n+1}$.*

- (1) *(X, x) is purely elliptic if and only if $(1, 1, \dots, 1) \in \Gamma(f)$.*
- (2) *Let $n = 3$ and let Δ_0 be the face of $\Gamma(f)$ consisting the point $(1, 1, 1, 1)$ in the relative interior of Δ_0 . Then we have*
 - (i) *(X, x) is a singularity of $(0, 2)$ -type if and only if $\dim_{\mathbb{R}} \Delta_0 = 3$.*
 - (ii) *(X, x) is a singularity of $(0, 1)$ -type if and only if $\dim_{\mathbb{R}} \Delta_0 = 2$.*
 - (iii) *(X, x) is a singularity of $(0, 0)$ -type if and only if $\dim_{\mathbb{R}} \Delta_0 = 1$ or $\dim_{\mathbb{R}} \Delta_0 = 0$.*

3 Principal parts

In this section, we give examples of the principal parts of hypersurface purely elliptic singularities of $(0, i)$ -type defined by a nondegenerate polynomial for $i = 0$ to $i = 1$.

EXAMPLE. Let (X, x) be the hypersurface purely elliptic singularity

$$xyzw + x^{5+p} + y^{5+q} + z^{5+r} + w^{5+s} = 0$$

in \mathbb{C}^4 . Blow up the point $O = (0, 0, 0, 0)$, let F be the exceptional set, and let Y be the strict transform of X . In this case the morphism $\pi : Y \rightarrow X$ is the canonical resolution of X . The exceptional set E consists of four 2-dimensional projective spaces in F , forming a tetrahedron.

EXAMPLE. Let (X, x) be the hypersurface purely elliptic singularity

$$x^2 + y^3 + z^7 + w^{43+s} + xyzw = 0$$

in \mathbb{C}^4 . Blow up the point $O = (0, 0, 0, 0)$ with weight $(21, 14, 6, 1)$, let F be the exceptional set, and let Y be the strict transform of X . In this case the morphism $\pi : Y \rightarrow X$ is the canonical resolution of X . The exceptional set E is a rational surface with a singularity $T_{2,3,7}$ in a weighted projective space F , i.e., $\mathbf{P}(21, 14, 6, 1)$.

EXAMPLE. Let (X, x) be the hypersurface singularity defined by the equation

$$x^2 + y^3 + z^7 + z^6 w^6 + w^{43+s} + xyzw = 0$$

in \mathbb{C}^4 . Then the singularity (X, x) is a purely elliptic singularity of $(0, 1)$ -type.

EXAMPLE. Let (X, x) be the hypersurface singularity defined by the equation

$$x^2 + y^3 + z^7 + \lambda z^6 w^6 + \mu w^{42} + w^{43+s} + xyzw = 0$$

in \mathbb{C}^4 . Then we obtain:

- (1) $\mu \neq 0 \Leftrightarrow (0, 2)$ -type.
- (2) $\mu = 0, \lambda \neq 0 \Leftrightarrow (0, 1)$ -type.
- (3) $\mu = 0, \lambda = 0 \Leftrightarrow (0, 0)$ -type.

References

- [1] S. Ishii, On isolated Gorenstein singularities, Math. Ann. 270 (1985), 541-554.
- [2] S. Ishii, The asymptotic behavior of plurigeners for a normal isolated singularity, Math Ann. 286 (1990), 803-812.
- [3] S. Ishii and K. Watanabe, On simple K3 singularities (in Japanese), Notes appearing in the Proceedings of the Conference on Algebraic Geometry at Tokyo Metropolitan Univ. (1988), 20-31.

- [4] S. Ishii and K. Watanabe, A geometric characterization of a simple K3 singularity, *Tohoku Math. J.* 44 (1992), 19-24.
- [5] A. N. Varchenko, Zeta-Function of monodromy and Newton's diagram, *Invent. Math.* 37 (1976), 253-262.
- [6] K. Watanabe, On plurigenera of normal isolated singularities, I, *Math. Ann.* 270 (1980), 65-94.
- [7] K. Watanabe, On plurigenera of normal isolated singularities, II, in *Complex Analytic singularities* (T. Suwa and P. Wagreich, eds.), *Advanced Studies in Pure Math.* 8, Kinokuniya, Tokyo and North-Holland. Amsterdam, New York, Oxford, (1986), 671-685.